

ERRATUM

Open Access



Erratum to: Analgesic use in a Norwegian general population: change over time and high-risk use - The Tromsø Study

Per-Jostein Samuelsen^{1,2*}, Lars Slørdal^{3,4}, Ulla Dorte Mathisen⁵ and Anne Elise Eggen²

Erratum

Upon publication of the original article [1], it was noticed that the following reference was erroneously left out:

Ademi Z, Turunen JH, Kauhanen J, Enlund H (2007) A comparison of three questionnaire-based measures of analgesic use over 11 years in adult males: a retrospective analysis of data from a prospective, population-based cohort study. *Clin Ther* 29 (3): 529–534

The reference should have been included after the sentence: “A comprehensive estimate of the use of analgesics cannot be done without the use of interview or questionnaires.” found towards the end of the Discussion section on page eight, second column, paragraph six, second sentence.

Author details

¹Regional Medicines Information and Pharmacovigilance Centre (RELIS), University Hospital of North Norway, N-9038 Tromsø, Norway. ²Department of Community Medicine, UiT-The Arctic University of Norway, Tromsø, Norway. ³Department of Laboratory Medicine, Children’s and Women’s Health, Norwegian University of Science and Technology, Trondheim, Norway. ⁴Department of Clinical Pharmacology, St. Olav University Hospital, Trondheim, Norway. ⁵Section of Nephrology, University Hospital of North Norway, Tromsø, Norway.

Received: 15 June 2017 Accepted: 20 June 2017

Published online: 27 June 2017

Reference

1. Samuelsen P-J, et al. Analgesic use in a Norwegian general population: change over time and high-risk use - The Tromsø Study. *BMC Pharmacol Toxicol.* 2015;16:16. doi:10.1186/s40360-015-0016-y.

* Correspondence: per-jostein.samuelsen@unn.no

¹Regional Medicines Information and Pharmacovigilance Centre (RELIS), University Hospital of North Norway, N-9038 Tromsø, Norway

²Department of Community Medicine, UiT-The Arctic University of Norway, Tromsø, Norway

