

POSTER PRESENTATION

Open Access

Identification and quantitation of 2',3'-cGMP in murine tissues

Heike Burhenne^{1,2*}, Sarah Tschirner¹, Roland Seifert¹, Volkhard Kaever^{1,2}

From 6th International Conference on cGMP: Generators, Effectors and Therapeutic Implications
Erfurt, Germany. 28-30 June 2013


Background

3',5'-cGMP is a well-known second messenger in eukaryotic cells which is synthesized by particulate and soluble guanylyl cyclases and is involved in several cardiovascular and neurological processes [1,2]. It has been recently reported that also 2',3'-cGMP (Figure 1) can be found in high concentrations in rabbit kidney and pancreas [3]. So far, little is known about the source and biological role of 2',3'-cGMP as well as its tissue distribution. To address this question, we have developed sensitive and specific liquid chromatography-coupled mass spectrometry (LC-MS/MS) methods for identification and quantitation of

2',3'- nucleoside monophosphates (2',3'-cNMPs) simultaneously with 3',5'-cNMPs (cAMP, cCMP, cGMP, cUMP). We systematically analyzed mouse tissues (brain, thymus, heart, lung, liver, pancreas, spleen, kidney, bladder, testis, ovary, uterus) for the presence of 2',3'- and 3',5'-cNMPs, respectively.


Materials and methods

For cNMP extraction, 30-200 mg tissue was transferred to a 2.0 mL tube containing a garnet matrix and one ¼ inch ceramic sphere. An organic extraction solvent was added and tissues were homogenated in a FastPrep-24[®] system



* Correspondence: burhenne.heike@mh-hannover.de

¹Institute of Pharmacology, Hannover Medical School, Germany
Full list of author information is available at the end of the article



(MP Biomedicals, Germany). The homogenate was centrifuged and the supernatant fluid was evaporated to dryness at 40°C under a nitrogen stream. The residue pellet was resuspended in water and analyzed by LC-MS/MS.

Detection and quantitation of 2',3'- and 3',5'-cNMPs was performed on a tandem mass spectrometer (5500 QTRAP®; AB SCIEX, USA). We confirmed our data on a TripleTOF™ 5600 system (AB SCIEX, USA) which is characterized by an extremely high mass accuracy.

Results

The chromatogram of a standard cNMP sample demonstrates that the LC-MS/MS method is suitable for the detection and quantitation of 2',3'- and 3',5'-cNMPs. Due to their retention times, all 2',3'-cNMPs could be reliably discriminated from their 3',5'-isomers (Figure 2A).

In addition to 3',5'-cGMP we detected notable amounts of 2',3'-cGMP in various mouse tissues. For example, heart samples showed an up to 5-fold higher concentration of 2',3'-cGMP compared with the 3',5'-cNMP (Figure 2B). In murine pancreas and spleen only 2',3'-cGMP but no 3',5'-cGMP could be detected.

Besides 2',3'-cGMP, remarkably high levels of 2',3'-cCMP and 2',3'-cUMP were detected in murine heart, kidney, spleen, liver, pancreas and lung.

Conclusion

2',3'-cAMP is an mRNA degradation product. Increased 2',3'-cAMP concentrations may play a role in cell death and cell proliferation [4].

We have demonstrated that high levels of 2',3'-cGMP as well as 2',3'-cCMP and 2',3'-cUMP can be detected in various murine tissues. Our results indicate that these cNMPs play an important but still unknown role in (patho)physiological processes.

To our knowledge, this is the first time that mammalian tissues were systematically analyzed for the occurrence of

2',3'-cNMPs. Our methods allow reliable detection and quantitation of four 2',3'-cNMPs simultaneously with their 3',5'-isomers and are, therefore, useful for the characterization of the physiological role of 2',3'-cNMPs.

Authors' details

¹Institute of Pharmacology, Hannover Medical School, Germany. ²Research Core Unit Mass Spectrometry - Metabolomics, Hannover Medical School, Germany.

Published: 29 August 2013

References

1. Kots AY, Martin E, Sharina IG, Murad F: A short history of cGMP, guanylyl cyclases, and cGMP-dependent protein kinases. *Handb Exp Pharmacol* 2009, **191**:1-14.
2. Stasch JP, Pacher P, Evgenov OV: Soluble guanylate cyclases as an emerging therapeutic target in cardiopulmonary disease. *Circulation* 2011, **123**:2263-2273.
3. Van Damme T, Zhang Y, Lynen F, Sandra P: Determination of cyclic guanosine- and cyclic adenosine monophosphate (cGMP and cAMP) in human plasma and animal tissues by solid phase extraction on silica and liquid chromatography-triple quadrupole mass spectrometry. *J Chromatogr B* 2012, **909**:14-21.
4. Jackson EK: The 2',3'-cAMP-adenosine pathway. *Am J Physiol Renal Physiol* 2011, **301**:F1160-F1167.

doi:10.1186/2050-6511-14-S1-P12

Cite this article as: Burhenne et al.: Identification and quantitation of 2',3'-cGMP in murine tissues. *BMC Pharmacology and Toxicology* 2013 **14** (Suppl 1):P12.