

POSTER PRESENTATION

Open Access

Absorption behavior of riociguat: bioavailability, food effects, and dose-proportionality

Corina Becker^{1*†}, Reiner Frey¹, Christiane Hesse¹, Sigrun Unger², Michael Reber¹, Wolfgang Mueck¹

From 6th International Conference on cGMP: Generators, Effectors and Therapeutic Implications Erfurt, Germany. 28-30 June 2013

Background

Riociguat, an oral soluble guanylate cyclase (sGC) stimulator, is currently being investigated for the treatment of pulmonary hypertension. Riociguat has a novel dual mode of action, directly stimulating sGC, independent of nitric oxide (NO), and increasing sensitivity of

sGC to NO. Riociguat thereby restores the NO-sGC-cGMP pathway, which is impaired in pulmonary hypertension. Three pharmacokinetic studies were performed to characterize the absorption behavior of riociguat including absolute bioavailability, food effects, and dose-proportionality.




Figure 1 Plasma riociguat concentrations following single 1 mg intravenous (n=22) or oral doses of riociguat (n=22) (absolute bioavailability study).

* Correspondence: corina.becker@bayer.com

† Contributed equally

¹Clinical Pharmacology, Bayer HealthCare Pharmaceuticals, Wuppertal, Germany

Full list of author information is available at the end of the article

Methods

The pharmacokinetic and safety profiles of riociguat were investigated in three open-label, randomized, crossover studies in healthy male subjects. In the absolute bioavailability study, fasted subjects ($n = 22$) received a single oral immediate-release dose of riociguat 1 mg or intravenous riociguat 1 mg. In the food-effects study, subjects ($n = 23$) received a single oral dose of immediate-release riociguat 2.5 mg after a 10-hour fast or after a high-fat high-calorie breakfast eaten within 30 minutes of dosing. In the dose-proportionality study, fasted subjects ($n = 24$) received a single oral dose of riociguat 0.5, 1, 1.5, 2, or 2.5 mg.

Results

In the absolute bioavailability study, riociguat exposure was similar after oral and intravenous dosing and oral bioavailability was 94% (95% confidence interval: 83–107)

(Figure 1). Mean C_{max} of riociguat was slightly lower after oral dosing compared with intravenous dosing. In the food-effects study, a high-fat breakfast had little effect on the extent of riociguat absorption, although absorption was delayed (Table 1). In fed subjects, the C_{max} of riociguat decreased by 35.3% and t_{max} increased compared with fasted subjects. In the dose-proportionality study, systemic exposure of riociguat was dose-proportional over 0.5–2.5 mg (Figure 2) with low intra-individual variability and moderate-to-high inter-individual variability. Riociguat was well tolerated in all studies. The most common treatment-emergent and riociguat-related adverse events were headache, flushing, and nasal congestion across all studies.

Conclusion

Riociguat shows complete oral absorption with no clinically relevant food effects; riociguat can therefore be

Table 1 Riociguat pharmacokinetics following a single oral dose of riociguat 2.5 mg in fed and fasted subjects (food-effects study)

Parameter ^a	Fasted (n=23)	Fed (n=23)	Estimated fed:fasted ratio (%)	90% CI	%CV
C_{max} ($\mu\text{g/L}$)	84.2 (44.7–152.7)	54.8 (28.9–91.4)	64.7	57.8–72.5	22.5
t_{max} (h)	1 (0.5–4.0)	4 (1.5–6.0)	—	—	—
$AUC_{0-\infty}$ ($\mu\text{g}\cdot\text{h/L}$)	572.2 (112.3–1300.0)	505.6 (113.3–1205.0)	88.3	82.2–95.0	14.3

AUC, area under the plasma concentration–time curve; CI, confidence interval; C_{max} , maximum riociguat plasma concentration; CV, coefficient of variation; t_{max} , time to reach C_{max} .

^aData are mean (range) except t_{max} , which is median (range).




Figure 2 Mean plasma riociguat concentrations following single oral doses of riociguat 0.5, 1.0, 1.5, 2.0, or 2.5 mg ($n=24$) (dose-proportionality study). LLOQ, lower limit of quantification.

taken with or without food. Riociguat systemic exposure increased dose proportionally over all doses (0.5–2.5 mg), supporting the suitability of the individualized dose-titration scheme used in the Phase III pulmonary arterial hypertension (PATENT) and chronic thromboembolic pulmonary hypertension (CHEST) studies.

Acknowledgements

The studies were funded by Bayer HealthCare Pharmaceuticals, Wuppertal, Germany. Medical writing assistance was provided by Adelphi Communications Ltd, Bollington, UK and funded by Bayer HealthCare Pharmaceuticals.

Authors' details

¹Clinical Pharmacology, Bayer HealthCare Pharmaceuticals, Wuppertal, Germany. ²Global Biostatistics, Bayer HealthCare Pharmaceuticals, Wuppertal, Germany.

Published: 29 August 2013

doi:10.1186/2050-6511-14-S1-P7

Cite this article as: Becker *et al.*: Absorption behavior of riociguat: bioavailability, food effects, and dose-proportionality. *BMC Pharmacology and Toxicology* 2013 **14**(Suppl 1):P7.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

