

MEETING ABSTRACT

Open Access

# Dephosphorylation of juxtamembrane serines and threonines of the NPR2 guanylyl cyclase regulates oocyte meiotic resumption

Leia C Shuhaibar<sup>1</sup>, Aaron B Edmund<sup>2</sup>, Jeremy R Egbert<sup>1</sup>, Siu-Pok Yee<sup>3</sup>, Lincoln R Potter<sup>2</sup>, Laurinda A Jaffe<sup>1\*</sup>


From 7th International Conference on cGMP Generators, Effectors and Therapeutic Implications  
Trier, Germany. 19-21 June 2015

## Background

The meiotic cell cycle of mammalian oocytes starts during embryogenesis and then pauses until luteinizing hormone (LH) restarts the cycle. This meiotic arrest is maintained by cGMP, which is produced in the granulosa cells by C-type natriuretic peptide (CNP) activation of NPR2 [1]. LH decreases cGMP in the granulosa cells, and via equilibration through gap junctions, cyclic GMP also decreases in the oocyte, thus releasing the meiotic arrest [2]. LH causes dephosphorylation and inactivation of NPR2 [3,4], but whether NPR2 dephosphorylation is required for meiotic resumption is not known. Seven regulatory NPR2 phosphorylation sites have been identified (Fig. 1) [5,6]. Here, we generated a knock-in mouse where each site was mutated to glutamate (*Npr2-7E*), resulting in a “constitutively phosphorylated” enzyme that we used to investigate the role of NPR2 dephosphorylation in the rapid resumption of meiosis in response to LH.

## Results


Membranes from isolated antral follicles treated with or without LH for 20 min were assayed for guanylyl cyclase activity under physiological conditions with CNP, ATP and Mg<sup>2+</sup>+GTP (Fig. 2). In wild-type follicles treated with LH, the CNP-dependent guanylyl cyclase activity decreased to ~47% of initial values (Fig. 2, left). In contrast, LH caused no significant change in CNP-dependent guanylyl cyclase activity in *Npr2-7E/7E* follicle membranes (Fig. 2, right). Protein loss did not explain the reduction in CNP-dependent activity because activities measured with



\* Correspondence: ljaffe@uchc.edu

<sup>1</sup>Department of Cell Biology, University of Connecticut Health Center, Farmington, Connecticut, USA

Full list of author information is available at the end of the article



**Figure 2** Inhibition of the LH-induced decrease in guanylyl cyclase activity in mouse follicles by the *Npr2* -7E mutations. Guanylyl cyclase activity in wild-type and *Npr2*-7E/7E follicle membranes, measured in the presence of 1  $\mu$ M CNP. Membranes were prepared from follicles with or without treatment with LH for 20 min. Each value shows the mean  $\pm$  s.e.m. for 4 separate membrane preparations. The asterisk indicates that LH treatment significantly decreased the CNP-dependent guanylyl cyclase activity in wild-type follicles ( $P < 0.05$ ). LH treatment did not cause a significant change in CNP-dependent activity in *Npr2*-7E/7E follicles (n.s.).

Mn<sup>2+</sup>+GTP/Triton X-100 were not reduced by LH. These findings indicate that dephosphorylation of NPR2 is necessary for the hormonal regulation of guanylyl cyclase activity in the ovarian follicle.

To investigate whether the LH-induced decrease in guanylyl cyclase activity is required for meiotic resumption, we isolated *Npr2*-7E/7E and wild-type follicles and observed them in culture before and after addition of LH. In wild-type follicles, LH-induced nuclear envelope breakdown (NEBD) began at 2 hours and reached ~80% by 6 hours. In contrast, in *Npr2*-7E/7E follicles, no evidence of meiotic resumption was observed in the first 6 hours following treatment with LH. However, by 8 hours after LH application, NEBD had occurred in ~40% of *Npr2*-7E/7E follicle-enclosed oocytes, and by 12 hours NEBD was seen in ~80% of oocytes. During cumulus expansion preceding ovulation, gap junction communication between the oocyte and cumulus cells is disrupted [7]. The LH-induced cumulus expansion occurred similarly in *Npr2*-7E/7E and wild-type follicles, suggesting that gap junction disruption during cumulus expansion might explain why meiosis eventually resumes in the NPR2-7E/7E follicles.

## Conclusion

Dephosphorylation of NPR2 is required for the rapid resumption of meiosis in response to LH.

## Acknowledgements

This work was supported by NIH grants R37HD014939, R01GM098309, T32AR050938 and the Fund for Science.

## Authors' details

<sup>1</sup>Department of Cell Biology, University of Connecticut Health Center, Farmington, Connecticut, USA. <sup>2</sup>Department of Biochemistry, Molecular Biology and Biophysics, University of Minnesota, Minneapolis, Minnesota, USA. <sup>3</sup>Department of Genetics and Genome Sciences, University of Connecticut Health Center, 263 Farmington Avenue, Farmington, CT 06030, USA.

Published: 2 September 2015

## References

1. Zhang M, Su Y-Q, Sugijara K, Xia G, Eppig JJ: Granulosa cell ligand NPPC and its receptor NPR2 maintain meiotic arrest in mouse oocytes. *Science* 2010, **330**:366-369.
2. Shuhaibar LC, Egbert JR, Norris RP, Lampe PD, Nikolaev VO, Thunemann M, Wen L, Feil R, Jaffe LA: Intercellular signaling via cyclic GMP diffusion through gap junctions in the mouse ovarian follicle. *Proc. Natl. Acad. Sci. U. S. A* 2015, in press.
3. Robinson JW, Zhang M, Shuhaibar LC, Norris RP, Geerts A, Wunder F, Eppig JJ, Potter LR, Jaffe LA: Luteinizing hormone reduces the activity of the NPR2 guanylyl cyclase in mouse ovarian follicles, contributing to the cyclic GMP decrease that promotes resumption of meiosis in oocytes. *Dev Biol* 2012, **366**:308-316.
4. Egbert JR, Shuhaibar LC, Edmund AB, Van Helden DA, Robinson JW, Uliasz TF, Baena V, Geerts A, Wunder F, Potter LR, Jaffe LA: Dephosphorylation and inactivation of the NPR2 guanylyl cyclase in the granulosa cells contributes to the LH-induced cGMP decrease that causes resumption of meiosis in rat oocytes. *Development* 2014, **141**:3594-3604.
5. Yoder AR, Stone MD, Griffin TJ, Potter LR: Mass spectrometric identification of phosphorylation sites in guanylyl cyclase A and B. *Biochemistry* 2010, **49**:10137-10145.
6. Yoder AR, Robinson JW, Dickey DM, Andersland J, Rose BA, Stone MD, Griffin TJ, Potter LR: A functional screen provides evidence for a conserved, regulatory, juxtamembrane phosphorylation site in guanylyl cyclase A and B. *PLoS One* 2012, **7**:e36747.
7. Gilula NB, Epstein ML, Beers WH: Cell-to-cell communication and ovulation. A study of the cumulus-oocyte complex. *J Cell Biol* 1978, **78**:58-75.

doi:10.1186/2050-6511-16-S1-A30

**Cite this article as:** Shuhaibar et al.: Dephosphorylation of juxtamembrane serines and threonines of the NPR2 guanylyl cyclase regulates oocyte meiotic resumption. *BMC Pharmacology and Toxicology* 2015 **16**(Suppl 1):A30.